

On the Orbit Decomposition of Finite Quandles

Sam Nelson

knots@esotericka.org

Chau-Yim Wong

jason@math.ucr.edu

Department of Mathematics, University of California, Riverside,
900 University Avenue, Riverside, CA, 92521

Abstract

We study the structure of finite quandles in terms of subquandles. Every finite quandle Q decomposes in a natural way as a union of disjoint Q -complemented subquandles; this decomposition coincides with the usual orbit decomposition of Q . Conversely, the structure of a finite quandle with a given orbit decomposition is determined by its structure maps. We describe a procedure for finding all non-connected quandle structures on a disjoint union of subquandles.

KEYWORDS: finite quandles, subquandles
2000 MSC: 57M27

1 Introduction

A *quandle* is a set Q with a binary operation $\triangleright$ satisfying

- (i) $x \triangleright x = x$ for every $x \in Q$,
- (ii) For every pair $x, y \in Q$, there is a unique $z \in Q$ such that $x = z \triangleright y$, and
- (iii) For every $x, y, z \in Q$, we have $(x \triangleright y) \triangleright z = (x \triangleright z) \triangleright (y \triangleright z)$.

If $(Q, \triangleright)$ satisfies (ii) and (iii), Q is a *rack*. Quandles form a category with morphisms $\phi : Q \rightarrow Q'$ defined as maps which preserve the quandle operation, i.e.

$$\phi(q_1 \triangleright q_2) = \phi(q_1) \triangleright' \phi(q_2)$$

where $\triangleright$ is the quandle operation in Q and $\triangleright'$ is the quandle operation in Q' . A bijective quandle homomorphism is a quandle isomorphism, as expected. Axiom (ii) implies that the map $f_b : Q \rightarrow Q$ defined by $f_b(a) = a \triangleright b$ is bijective for all $b \in Q$; the inverse then defines a second operation $a \triangleright^{-1} b = f_b^{-1}(a)$, called the *dual* operation of Q ; the quandle $(Q, \triangleright^{-1})$ is the *dual* of Q . Quandles have been studied in many recent papers; see [6] for more on the history of racks and quandles.

Standard examples of quandles include Alexander quandles, i.e., modules M over the ring $\Lambda = \mathbb{Z}[t^{\pm 1}]$ of Laurent polynomials in one variable with integer coefficients with quandle operation given by

$$x \triangleright y = tx + (1 - t)y,$$

and groups, which are quandles with quandle operation given by $x \triangleright y = y^{-1}xy$. If the group is abelian, the quandle operation reduces to $x \triangleright y = x$, and the quandle is *trivial*.

Quandles are of interest to topologists since the knot quandle (see [10]) is a complete invariant of knots up to homeomorphism of topological pairs. Finite quandles are of particular interest as a source of computable knot invariants such as the counting invariants $|\text{Hom}(K, C)|$ where K is a knot quandle and C is a finite coloring quandle as well as related invariants which make use of various quandle cohomology theories (see [4], [3], etc.)

In this paper, we study the structure of finite quandles in terms of subquandles. Our initial goal was to try to find something like a Sylow theorem for finite quandles. In section 2, we study the structure of finite quandles in terms of orbit subquandles. We show how to determine all quandles with a given two-subquandle orbit decomposition and we discuss how to find quandle structures on a union of three or more orbit subquandles.

In section 3, we use quandle matrices (see [9]) to study the structure of finite quandles. We describe algorithms for finding the orbit decomposition of a finite quandle and for finding quandle structures on a disjoint union of n subquandles. This is intended to lay the groundwork for the related problem of counting the number of ways of filling in zeroes in a quandle presentation matrix to obtain a finite quandle; it is hoped that a solution to this problem might give new insights into the quandle-counting invariants of knots and links studied in various recent papers ([3], [5]). *Maple* code for finding rack actions and orbit decompositions is available at www.esotericka.org/quandles.

2 Orbit Decomposition

Let Q be a quandle. A *subquandle* $X \subset Q$ is a subset of Q which is itself a quandle under $\triangleright$. Unlike the case of groups, in which the intersection of any collection of subgroups is always non-empty (containing at least the identity element), a collection of subquandles of a given quandle may be pairwise disjoint. Indeed, unlike groups, every subset $X \subset Q$ which is closed under $\triangleright$ is a subquandle: if X is closed under $\triangleright$, the restriction $f_b|_X : X \rightarrow X$ is bijective for every $b \in X$, and axiom (ii) is satisfied. Since axioms (i) and (iii) are automatic for any subset of Q , this makes X a subquandle. Thus we have

Lemma 1 *Let $(Q, \triangleright)$ be a quandle and $X \subset Q$ a subset. Then X is a subquandle iff X is closed under $\triangleright$.*

We would like to understand the structure of a quandle in terms of its subquandles. A quandle which can be written as a union of two disjoint subquandles has been called *decomposable* in the literature (see [1], [7], [11], etc.), and a quandle which is not a disjoint union of two subquandles is *indecomposable*. The existence of indecomposable quandles follows from the fact that the complement of a subquandle is not necessarily a subquandle. However, as observed in [11], indecomposability of a quandle Q does not imply that Q has no subquandles, nor even that the quandle cannot be decomposed as a disjoint union of three or more subquandles.

Indeed, every singleton subset of a quandle is itself a subquandle, though the analogous statement is not true for non-quandle racks. Thus every quandle decomposes in an unhelpful way as a disjoint union of singleton subquandles. In [11], we find the dihedral quandle R_9 , an indecomposable quandle which can be written as a disjoint union of three isomorphic subquandles. Though this quandle is “indecomposable,” since the complements of each of the three subquandles are not closed under $\triangleright$, it nevertheless has an internal structure determined by its component subquandles – namely, a Cartesian product of a quandle of order three with itself. This is an example of a congruence structure (see [12]).

Definition 1 Let Q be a quandle and $X \subset Q$ a subquandle. We say that X is *complemented in Q* or *Q -complemented* if $Q \setminus X$ is a subquandle of Q . Note that since the empty set $\emptyset$ is a quandle, every quandle Q is complemented in itself. A quandle Q is *complementary* if it has a nonempty Q -complemented subquandle.

A complementary quandle Q may have subquandles which are not Q -complemented; indeed, every singleton subset of Q is a subquandle, while in general $Q \setminus \{x\}$ is not a subquandle. It is clear from the definition of decomposability that a quandle is decomposable iff it is complementary.

[10] includes the following definition:

Definition 2 A quandle is *algebraically connected* or just *connected* if for every $a, b \in Q$, we have

$$(\dots((a \diamond_1 x_1) \diamond_2 x_2) \dots) \diamond_n x_n = b$$

for some $x_1, \dots, x_n \in Q$, $\diamond_1 \dots \diamond_n \in \{\triangleright, \triangleright^{-1}\}$. The set of all such $b \in Q$ is the *orbit* of a .

It is well-known that algebraic connectedness coincides with indecomposability in the sense defined above, and hence coincides with non-complementarity.

Lemma 2 Let X and Y be subquandles of a quandle Q . Then $X \cap Y$ is a subquandle. If X and Y are Q -complemented, so is $X \cap Y$.

Proof. Let $x, y \in X \cap Y$. Then $x \triangleright y \in X$ since X is a subquandle, and $x \triangleright y \in Y$ since Y is a subquandle. Hence $x \triangleright y \in X \cap Y$, and $X \cap Y$ is closed under $\triangleright$, and $X \cap Y$ is a subquandle by lemma 1.

Now, suppose X and Y are Q -complemented; we must show that $Z = Q \setminus (X \cap Y) = (Q \setminus X) \cup (Q \setminus Y)$ is a subquandle. Let $x, y \in Z$. If x and y are both in $Q \setminus X$ or both in $Q \setminus Y$ then $x \triangleright y \in Z$ since $Q \setminus X$ and $Q \setminus Y$ are closed under $\triangleright$. If $x \in Q \setminus X$ and $y \notin Q \setminus X$, then $y \in X$, which implies $w = x \triangleright y \in Q \setminus X \subset Z$, since otherwise the closure of X under $\triangleright^{-1}$ would imply $w \triangleright^{-1} y = x \in X$, contradicting our choice of x . Similarly, $y \in Q \setminus Y$ and $x \in Z \setminus (Q \setminus X)$ implies $y \triangleright x \in Q \setminus Y \subset Z$, and Z is closed under $\triangleright$ as required. $\square$

Theorem 3 Let Q be a finite quandle. Then Q may be written as

$$Q = Q_1 \amalg Q_2 \amalg \dots \amalg Q_n$$

where every Q_i is Q -complemented and no proper subquandle of any Q_i is Q -complemented. This decomposition is well-defined up to isomorphism; if $Q \cong Q'$, then in the decompositions

$$Q = Q_1 \amalg \dots \amalg Q_n \quad \text{and} \quad Q' = Q'_1 \amalg \dots \amalg Q'_m,$$

we have then $n = m$ and (after reordering if necessary), $Q_i \cong Q'_i$.

Remark 1 The decomposition of a finite quandle into orbits coincides with our notion of decomposition into Q -complemented subquandles; this follows from the observation that the orbits in Q are Q -complemented subquandles. Q -complemented subquandle decomposition then gives us a new perspective on the division of Q into disjoint orbits. Indeed, we will see how to construct a quandle with a specified list of orbits, when such exists. See also [1] proposition 1.17.

Proof. For every $a \in Q$, define $S(a)$ to be the intersection of all Q -complemented subquandles of Q containing a . The collection $\{S(a) \mid a \in Q\}$ is the orbit decomposition of Q : each $S(a)$ is Q -complemented, no proper subquandle of any $S(a)$ is Q -complemented, and $S(a) \cap S(b) \neq \emptyset$ implies $S(a) = S(b)$. Since the empty quandle is Q -complemented, if Q has no nonempty Q -complemented proper subquandles, then $\{S(a) \mid a \in Q\} = \{Q\}$; in any case, $\cup_{a \in Q} S(a) = Q$.

If $\phi : Q \rightarrow Q'$ is an isomorphism, then for any subquandle $S \subset Q$ the restriction $\phi|_S$ is an isomorphism onto a subquandle of Q' . In particular, if S is Q -complemented, then $\phi|_{Q \setminus S}$ is also an isomorphism onto the subquandle $Q' \setminus \phi(S')$. Hence Q' has an isomorphic list of Q' -complemented subquandles before taking intersections, and thus has an isomorphic orbit decomposition. $\square$

Example 1 Let Q be the trivial quandle $T_n = \{1, 2, \dots, n\}$ with quandle operation $i \triangleright j = i$ for all $i, j \in Q$. Then every singleton subquandle $\{i\} \subset Q$ is Q -complemented, so the orbit decomposition of T_n is the maximal partition $T_n = \{1\} \amalg \{2\} \amalg \dots \amalg \{n\}$.

Before we come to the next theorem, we need a definition.

Definition 3 Let R be a rack and S a quandle. A *rack action* of R on S is a map from R to the set of automorphisms of S , $\Phi : R \rightarrow \text{Aut}(S) = \{\phi_r : S \rightarrow S, \quad r \in R\}$, such that

$$\phi_r(\phi_{r'}(s)) = \phi_{r' \triangleright r}(\phi_r(s))$$

for all $r, r' \in R$ and for all $s \in S$.

Example 2 Let Q be a quandle. Then the set $F : Q \rightarrow \text{Aut}(Q) = \{f_y : Q \rightarrow Q \mid f_y(x) = x \triangleright y\}$ is a rack action of Q on itself, since

$$f_z(f_y(x)) = (x \triangleright y) \triangleright z = (x \triangleright z) \triangleright (y \triangleright z) = f_{y \triangleright z}(f_z(x))$$

for all $x, y, z \in Q$.

Remark 2 Lemma 4 is a special case of lemma 1.18 in [1].

Lemma 4 Let Q and Q' be finite quandles. Then there is a quandle $U = Q \amalg Q'$ iff there are rack actions $F : Q \rightarrow \text{Aut}(Q')$ and $G : Q' \rightarrow \text{Aut}(Q)$ such that the compatibility conditions

$$g_x(a) \triangleright b = g_{f_b(x)}(a \triangleright b) \quad \text{and} \quad f_a(x) \triangleright y = f_{g_y(a)}(x \triangleright y)$$

are satisfied for all $a, b \in Q, x, y \in Q'$.

Proof. ($\Rightarrow$) Let $U = Q \cup Q'$ and let F, G be rack actions. Define

$$x \triangleright y = \begin{cases} f_y(x) & x \in Q', y \in Q \\ g_y(x) & x \in Q, y \in Q'. \end{cases}$$

Then we assert that $(U, \triangleright)$ is a quandle. The first quandle axiom is satisfied because Q and Q' are quandles themselves. The second axiom follows from the definition of rack action: each f_a and g_x defines a bijection of Q' and Q respectively, while the fact that Q and Q' are quandles says that

each element acts as a bijection on its own subquandle. Hence the action of b on $U = Q \amalg Q'$ is bijective for each $b \in U$, and axiom (ii) is satisfied.

To see that U satisfies (iii), we simply check all the possibilities. If q_1, q_2 and q_3 are in the same subquandle, then (iii) is satisfied. If $q_1 \in Q$ and $q_2, q_3 \in Q'$, then

$$(q_1 \triangleright q_2) \triangleright q_3 = g_{q_3}(g_{q_2}(q_1)) = g_{q_2 \triangleright q_3}(g_{q_3}(q_1)) = (q_1 \triangleright q_3) \triangleright (q_2 \triangleright q_3)$$

since G is a rack action. Similarly, the fact that F is a rack action implies that $(q_1 \triangleright q_2) \triangleright q_3 = (q_1 \triangleright q_3) \triangleright (q_2 \triangleright q_3)$ when $q_1 \in Q'$ and $q_2, q_3 \in Q$.

If $q_1, q_2 \in Q$ and $q_3 \in Q'$, then

$$(q_1 \triangleright q_2) \triangleright q_3 = g_{q_3}(q_1 \triangleright q_2) = g_{q_3}(q_1) \triangleright g_{q_3}(q_2) = (q_1 \triangleright q_3) \triangleright (q_2 \triangleright q_3)$$

since g_{q_3} is quandle homomorphism for each $q_3 \in Q'$. Similarly $(q_1 \triangleright q_2) \triangleright q_3 = (q_1 \triangleright q_3) \triangleright (q_2 \triangleright q_3)$ when $q_1, q_2 \in Q'$ and $q_3 \in Q$ since each f_{q_3} is a quandle homomorphism.

Finally, if $q_1, q_3 \in Q'$ and $q_2 \in Q$, then

$$(q_1 \triangleright q_2) \triangleright q_3 = g_{q_3}(f_{q_2}(q_1)) = f_{g_{q_3}(q_2)}(q_1 \triangleright q_3) = (q_1 \triangleright q_3) \triangleright (q_2 \triangleright q_3).$$

Similarly, the compatibility condition ensures that $(q_1 \triangleright q_2) \triangleright q_3 = (q_1 \triangleright q_3) \triangleright (q_2 \triangleright q_3)$ when $q_1, q_3 \in Q$ and $q_2 \in Q'$.

($\Leftarrow$) If $U = Q \amalg Q'$, then one easily checks that $F : Q \rightarrow \text{Aut}(Q')$ and $G : Q' \rightarrow \text{Aut}(Q)$ are rack actions satisfying the compatibility condition. $\square$

Indeed, as noted in [1], lemma 4 can be easily generalized to obtain:

Theorem 5 *Let $Q_1, \dots, Q_n$ be finite quandles. Then there is a quandle $Q = \amalg_{i=1}^n Q_i$ if there are rack actions $\Phi^{i,j} : Q_i \rightarrow \text{Aut}(Q_j)$ satisfying the compatibility conditions*

$$\phi_c^{k,i}(\phi_b^{j,i}(a)) = \phi_{\phi_c^{k,j}(b)}^{k,i}(\phi_c^{k,i}(a))$$

for all $a \in Q_i$, $b \in Q_j$ and $c \in Q_k$. Moreover, $\{Q_1, \dots, Q_n\}$ is the orbit decomposition of Q unless all the rack actions $\Phi^{i,j}$ preserve a Q_j -complemented subquandle $A \subset Q_j$ for some Q_j .

Proof. As above, for each $x \in Q_i$, $y \in Q_j$ define $x \triangleright y = \phi_y^{j,i}(x)$. Then quandle axiom (i) is satisfied automatically since each Q_i is itself a quandle. Axiom (ii) is satisfied since each element acts on each of the disjoint subquandles Q_i by an automorphism, so the overall action is a bijection for each element.

Axiom (iii) is satisfied by the compatibility conditions when the three elements are in distinct subquandles or when a and c are in one subquandle and b is in another. For example, if $a \in Q_i$, $b \in Q_j$ and $c \in Q_k$ we have

$$(a \triangleright b) \triangleright c = \phi_b^{j,i}(a) \triangleright c = \phi_c^{k,i}(\phi_b^{j,i}(a)) = \phi_{\phi_c^{k,j}(b)}^{k,i}(\phi_c^{k,i}(a)) = \phi_{b \triangleright c}^{k,i}(a \triangleright c) = (a \triangleright c) \triangleright (b \triangleright c).$$

As before, the rack action and automorphism requirements satisfy axiom (iii) in the other cases.

Finally, note that removing any Q_i from the list along with the corresponding $\Phi^{i,j}$ and $\Phi^{j,i}$ rack actions still defines a quandle, so the Q_i s are each Q -complemented. If no subquandle of any Q_i is preserved by all the actions $\Phi^{i,j}$, then no subquandle of Q_i is Q -complemented and $\{Q_1, \dots, Q_n\}$ is the orbit decomposition of Q . $\square$

Definition 4 Call the maps $\Phi^{i,j}$ the *structure maps* of the quandle U with respect to the decomposition $U = Q_1 \amalg Q_2 \amalg \cdots \amalg Q_n$.

Corollary 6 *There is a quandle Q with orbits $Q_1, \dots, Q_n$ iff there are compatible rack actions $\Phi^{i,j} : Q_i \rightarrow \text{Aut}(Q_j)$ such that for every Q_j -complemented proper subquandle $S \subset Q_j$ at least one of the automorphisms $\Phi_a^{i,j}$ for some $a \in Q_i$ does not satisfy $\Phi_a^{i,j}(S) = S$.*

Proof. If $S \subset Q_j$ is a proper Q_j -complemented subquandle such that every $\Phi_a^{i,j}(S) = S$, then S is an orbit of Q . $\square$

Example 3 Let $Q_1, \dots, Q_n$ be any finite collection of finite quandles, and define $\Phi^{i,j} : Q_i \rightarrow \text{Aut}(Q_j)$ by $\phi_x^{i,j} = \text{Id}_{Q_j}$ for all $x \in Q_i$. Then

$$\phi_x^{i,j}(\phi_y^{i,j}(q)) = q = \phi_{y \triangleright x}^{i,j}(\phi_x^{i,j}(q))$$

for all $q \in Q_j$, so each $\Phi^{i,j}$ is a rack action. Moreover,

$$\phi_c^{k,i}(\phi_b^{j,i}(a)) = a = \phi_{\phi_c^{k,j}(b)}^{k,i}(\phi_c^{k,i}(a))$$

for all i, j, k , so $U = Q_1 \amalg \cdots \amalg Q_n$ is a quandle. This example shows that there is always at least one quandle structure on the union of any finite collection of finite quandles. Indeed, if the subquandles $Q_1, \dots, Q_n$ are non-complementary, then the orbits of U are precisely $Q_1, \dots, Q_n$. This example is sometimes called the *disjoint union* of the subquandles $Q_1, \dots, Q_n$ (see [12]).

Example 4 Let $Q_i = \{x_i\}, i = 1 \dots n$ be a collection of n singleton quandles. Then the only possible rack actions by automorphisms of singleton quandles on other singleton quandles are the identity actions, so the only quandle structure with orbit decomposition consisting of all singletons must have $x_i \triangleright x_j = x_i$ for all $x_i, x_j \in Q$, that is, the trivial quandle T_n .

The observation that a quandle $Q_j \subset Q$ may have subquandles which are Q_j -complemented but not Q -complemented implies that quandle may have multiple layers of orbit decompositions. Specifically, if $U = Q_1 \amalg \cdots \amalg Q_n$ is the orbit decomposition of U , then each Q_i will have its own orbit decomposition, consisting of multiple subquandles if Q_i is not connected. Define the *subquandle depth* of U to be the maximum number n of such layers of decomposition needed before all remaining orbit decompositions consist of connected subquandles. This subquandle depth is an invariant of quandle isomorphism type. A connected quandle has subquandle depth 0; indeed, we may take this as an alternate definition for “connected.”

Example 5 The quandle U with quandle matrix (see the next section)

$$M_U = \begin{bmatrix} 1 & 1 & 2 & 2 \\ 2 & 2 & 1 & 1 \\ 4 & 4 & 3 & 3 \\ 3 & 3 & 4 & 4 \end{bmatrix}$$

has orbit decomposition $U = \{1, 2\} \amalg \{3, 4\}$. The two orbit subquandles are trivial and thus have further orbit decompositions $\{1\} \amalg \{2\}$ and $\{3\} \amalg \{4\}$, so this quandle has subquandle depth 2.

The structure maps of theorem 5 define a quandle structure on $U = Q_1 \amalg \cdots \amalg Q_n$ with subquandle depth 1. To find all non-connected quandle structures on U with subquandles $Q_1, \dots, Q_n$ we must consider not only quandle structures with subquandle depth 1 but all other possible subquandle depths. To obtain the list of all subquandle depth 2 quandle structures on U we must consider all partitions the set $\{Q_1, \dots, Q_n\}$ into disjoint subsets. Then for each partition we apply theorem 5 to each subset in the partition, obtaining the list of all subquandle depth 1 quandles on each subset of the partition. Applying the theorem again to the resulting new lists of quandles yields a set including all subquandle depth 2 quandle structures on U . Applying this procedure recursively – that is, for each set S in a partition of $\{Q_1, \dots, Q_n\}$, consider all the partitions of S , etc. – yields all non-connected quandle structures on U such that each Q_i is a subquandle of U , since every non-connected quandle structure has some subquandle depth between 1 and n . Note that the sets of quandle structures obtained from distinct partitions of $\{Q_1, \dots, Q_n\}$ are not disjoint – the structure in which all rack actions are trivial, for example, can be built from any partition (or partitioned partition, etc.).

Summarizing, we have

Corollary 7 *Every non-connected quandle structure on $U = Q_1 \amalg \cdots \amalg Q_n$ with Q_i subquandles has an orbit decomposition recursively obtainable from quandles with orbit decompositions consisting of subsets of $Q_1 \amalg \cdots \amalg Q_n$.*

Finally, if $\rho : Q \rightarrow Q'$ is an isomorphism of quandles where $Q = Q_1 \amalg Q_2 \amalg \cdots \amalg Q'_n$ and $Q' = Q'_1 \amalg Q'_2 \amalg \cdots \amalg Q'_n$ are the orbit decompositions of Q and Q' , then denoting $\rho|_{Q_i} = \rho_i$, we have

$$\rho_j(\phi_b^{i,j}(a)) = \rho(a \triangleright b) = \rho(a) \triangleright \rho(b) = \phi_{\rho_i(b)}^{i,j}(\rho_j(a)).$$

Hence we have

Proposition 8 *Let $Q_1, \dots, Q_n$ be finite quandles. Then two quandle structures on the union $U = Q_1 \amalg \cdots \amalg Q_n$ given by $\Phi^{i,j}$ and $\Psi^{i,j}$ are isomorphic iff there are automorphisms $\rho_i : Q_i \rightarrow Q_i$ such that*

$$\rho_j(\phi_b^{i,j}(a)) = \psi_{\rho_i(b)}^{i,j}(\rho_j(a))$$

for all $a \in Q_j, b \in Q_i$.

3 Quandle Matrices and computation

The lists of quandles of order $n \leq 6$ in [2] and order $n \leq 5$ in [9] show that many of the possible quandle structures of small order may be understood as unions of disjoint subquandles. This observation naturally raises the question of how many different ways there are for two (or more) quandles to be put together, that is, how many quandle structures are possible on $Q \amalg Q'$ which have Q and Q' as subquandles. Theorem 5 gives us an answer, but we need some more convenient notation in order to permit computations. Quandle matrix notation (see [9]) provides a solution for this problem.

Definition 5 Let $Q = \{x_1, x_2, \dots, x_n\}$ be a quandle. The *matrix* of Q , M_Q , is the matrix abstracted from the operation table of Q by forgetting the x s and keeping only the subscripts. That is, we set $(M_Q)_{ij} = k$ where $x_k = x_i \triangleright x_j$ in Q .

Note that quandle axiom (i) enables us to deduce row and column labels and hence recover Q from M_Q .

Example 6 The Alexander quandle $Q_3 = \Lambda_3/(t+1) = \{x_1 = 0, x_2 = 1, x_3 = 2\}$ has operation table

$$\begin{array}{c|ccc} & x_1 & x_2 & x_3 \\ \hline x_1 & x_1 & x_3 & x_2 \\ x_2 & x_3 & x_2 & x_1 \\ x_3 & x_2 & x_1 & x_3 \end{array} \quad \text{and thus} \quad M_Q = \begin{bmatrix} 1 & 3 & 2 \\ 3 & 2 & 1 \\ 2 & 1 & 3 \end{bmatrix}.$$

A non-trivial quandle may have an orbit decomposition into trivial subquandles, and the orbits of a quandle need not be connected.

Example 7 The quandle defined by the quandle matrix

$$\begin{bmatrix} 1 & 1 & 1 & 2 \\ 2 & 2 & 2 & 3 \\ 3 & 3 & 3 & 1 \\ 4 & 4 & 4 & 4 \end{bmatrix}$$

has orbit decomposition as $T_3 \amalg T_1$ with rack actions $F : T_3 \rightarrow T_1$ given by $f_1(x) = f_2(x) = f_3(x) = x$ and $G : T_1 \rightarrow T_3$ given by $g_4(1) = 2, g_4(2) = 3$, and $g_4(3) = 1$, that is, g_4 is the permutation (123).

Just as matrix notation provides a convenient way to specify finite quandles, we can use column vector notation to represent maps from one finite quandle to another. Specifically, given a map $\phi : Q \rightarrow Q'$ where $Q = \{1, 2, \dots, n\}$ and $Q' = \{1, 2, \dots, m\}$ are quandles given by matrices M_Q and $M_{Q'}$, we can represent the map ϕ as the n -component column vector

$$\phi = \begin{bmatrix} \phi(1) \\ \phi(2) \\ \vdots \\ \phi(n) \end{bmatrix}, \quad \phi(i) \in \{1, \dots, m\}.$$

Then a rack action $F : Q \rightarrow \text{Aut}(Q')$ may be represented as an $m \times n$ matrix where the i th column is the vector representation of $f_i : Q' \rightarrow Q'$. Lemma 4 then gives us an algorithm for determining all quandle structures on $Q \amalg Q'$, namely let $Q = \{1, 2, \dots, n\}$ and $Q' = \{n+1, \dots, n+m\}$ be quandles with matrices M_Q and $M_{Q'}$ respectively. Then

- (1) For every $m \times n$ matrix F with columns which are permutations of Q' , check whether the matrix satisfies the rack action condition

$$F[i, j] = F[M_Q[i, j], i] \quad \forall i, j \in Q$$

- (2) For every $n \times m$ matrix G with columns which are permutations of Q , check whether the matrix satisfies the rack action condition

$$G[i, j] = G[M_{Q'}[i, j], i] \quad \forall i, j \in Q'$$

(3) For every pair F, G of such matrices, test the compatibility conditions

$$M[F[i, k], j] = G[M'[i, k], F[G[k, i]]] \quad \forall i, k \in Q, j \in Q'$$

and

$$M'[G[i, k], j] = F[M[i, k], G[F[k, i]]] \quad \forall i, k \in Q', j \in Q.$$

(4) For every pair F, G which passes steps (1)-(3), the block matrix

$$\left[\begin{array}{c|c} M_Q & G \\ \hline F & M'_Q \end{array} \right]$$

is a quandle matrix.

Conversely, given a quandle matrix Q_M , we can read off the rack actions by simply interpreting Q_M as a block matrix.

We note that the generalization of this procedure to unions of more than two quandles does not give all possible quandle structures on the disjoint union of three or more subquandles, since this construction yields only quandles in which every given Q_i is Q -complemented, i.e., quandle structures of subquandle depth 1.

For example, the connected quandle $Q_3 \times Q_3$ has matrix

$$M_{Q_3 \times Q_3} = \left[\begin{array}{ccc|ccc|ccc} 1 & 3 & 2 & 7 & 9 & 8 & 4 & 6 & 5 \\ 3 & 2 & 1 & 9 & 8 & 7 & 6 & 5 & 4 \\ 2 & 1 & 3 & 8 & 7 & 9 & 5 & 4 & 6 \\ \hline 7 & 9 & 8 & 4 & 6 & 5 & 1 & 3 & 2 \\ 9 & 8 & 7 & 6 & 5 & 4 & 3 & 2 & 1 \\ 8 & 7 & 9 & 5 & 4 & 6 & 2 & 1 & 3 \\ \hline 4 & 6 & 5 & 1 & 3 & 2 & 7 & 9 & 8 \\ 6 & 5 & 4 & 3 & 2 & 1 & 9 & 8 & 7 \\ 5 & 4 & 6 & 2 & 1 & 3 & 8 & 7 & 9 \end{array} \right],$$

which has no $Q_3 \times Q_3$ -complemented subquandles. This quandle is isomorphic to the dihedral quandle R_9 whose three-subquandle decomposition is noted in [11]. Indeed, we can use the division algorithm to write a quandle matrix for $Q \times Q'$ where $|Q| = n$ and $|Q'| = m$ by identifying (x, y) with $(x-1)m + y$ for $x = 1, \dots, n$ and $y = 1, \dots, m$. Then the matrix of $Q \times Q'$ is the block matrix

$$\left[\begin{array}{c|c|c|c} (q_{11} - 1)m + M_{Q'} & (q_{12} - 1)m + M_{Q'} & \dots & (q_{1m} - 1)m + M_{Q'} \\ \hline (q_{21} - 1)m + M_{Q'} & (q_{22} - 1)m + M_{Q'} & \dots & (q_{2m} - 1)m + M_{Q'} \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline (q_{m1} - 1)m + M_{Q'} & (q_{m2} - 1)m + M_{Q'} & \dots & (q_{mm} - 1)m + M_{Q'} \end{array} \right]$$

where $M_Q = (q_{ij})$ and $M_{Q'}$ are the matrices of Q and Q' respectively.

Remark 3 In the last section, we noted that if a finite quandle Q is a union of three or more subquandles, then some quandle structures may have subquandle depth greater than 1, since $x \in Q_1$ need not imply $x \triangleright y \in Q_1$. For example, the quandle Q below is a union of three subquandles $Q_1 = \{1, 2\}$, $Q_2 = \{3, 4\}$ and $Q_3 = \{5, 6\}$, and indeed there is an apparent block-matrix decomposition.

However, because Q_1 is not Q -complemented, there is no rack action $\Phi^{2,1} : Q_2 \rightarrow \text{Aut}(Q_1)$, for example.

$$M_Q = \left[\begin{array}{cc|cc|cc} 1 & 1 & 2 & 2 & 1 & 1 \\ 2 & 2 & 5 & 5 & 2 & 2 \\ \hline 3 & 3 & 3 & 3 & 3 & 3 \\ 4 & 4 & 4 & 4 & 4 & 4 \\ \hline 5 & 5 & 1 & 1 & 5 & 5 \\ 6 & 6 & 6 & 6 & 6 & 6 \end{array} \right]$$

To construct this subquandle depth 2 quandle from Q_1, Q_2 and Q_3 we must first put together $Q' = Q_1 \amalg Q_3 \cong T_4$, then find structure maps for $Q = Q' \amalg Q_2$.

Maple programs for finding rack actions and orbit decompositions of finite quandles represented by matrices are available in the file `quandles-maple.txt` at www.esotericka.org/quandles.

References

- [1] N. Andruskiewitsch and M. Graña. From racks to pointed Hopf algebras. *Adv. Math.* **178** (2003) 177-243.
- [2] J. S. Carter, S. Kamada and M. Saito. *Surfaces in 4-space*. Encyclopaedia of Mathematical Sciences, 142. Low-Dimensional Topology, III. Springer-Verlag, Berlin, 2004.
- [3] J. S. Carter, D. Jelsovsky, S. Kamada, L. Langford and M. Saito. Quandle cohomology and state-sum invariants of knotted curves and surfaces. *Trans. Amer. Math. Soc.* **355** (2003) 3947-3989.
- [4] J. S. Carter, M. Elhamdadi, M. Graña and M. Saito. Cocycle Knot Invariants from Quandle Modules and Generalized Quandle Cohomology. *Osaka J. Math.* **42** (2005) 499-541.
- [5] F. M. Dionísio and P. Lopes. Quandles at finite temperatures. II. *J. Knot Theory Ramifications* **12** (2003) 1041-1092.
- [6] R. Fenn and C. Rourke. Racks and links in codimension two. *J. Knot Theory Ramifications* **1** (1992) 343-406.
- [7] P. Etingof and M. Graña. On rack cohomology. *J. Pure Appl. Algebra* **177** (2003) 49-59.
- [8] R. Henderson, T. Macedo and S. Nelson Symbolic Computation with finite quandles. *J. Symbolic Comput.* **41** (2006) 811-817.
- [9] B. Ho and S. Nelson. Matrices and Finite Quandles. *Homology Homotopy Appl.* **7** (2005) 197-208.
- [10] D. Joyce. A classifying invariant of knots, the knot quandle. *J. Pure Appl. Algebra* **23** (1982) 37-65.
- [11] P. Lopes and D. Roseman. On finite racks and quandles. *Comm. Algebra* **34** (2006) 371-406.
- [12] H. Ryder. *The Structure of Racks*. Ph.D. Dissertation, U. Warwick. Available online at www.esotericka.org/quandles